






Identification of commercially available antibodies that block ligand-binding by BMPR2

Ruthann Gorrell & Jonathan W. Lowery, PhD

Division of Biomedical Science, Marian University College of Osteopathic Medicine, Indianapolis, Indiana, USA

Osteoporosis, a disease of low bone mineral density, affects 10 million Americans and is a significant health problem and a considerable socioeconomic burden. Current treatments for osteoporosis have significant limitations, necessitating identifying new treatment strategies via building a better understanding of the endogenous mechanisms regulating bone mass. Recent research demonstrated that removal of the BMP type 2 receptor, BMPR2, in skeletal progenitor cells of *Bmpr2*-cKO mice leads to reduced age-related bone loss due to a sustained elevation in bone formation rate. The molecular mechanism underlying this phenotype is being pursued in other work. In the present study, we sought to advance the translational potential of the genetic model by identifying antibodies that neutralize the ligand-binding function of the BMPR2 extracellular domain (BMPR2-ECD). Using a modified, cell-free immunoprecipitation assay quantified by ELISA, we examined the neutralizing ability of 3F6, which is a mouse monoclonal antibody raised against the ligand-binding region of BMPR2, and found a dose-dependent inhibition of BMPR2-ECD ligand-binding. We then evaluated 1F12, which is another mouse monoclonal antibody raised against the ligand-binding region of BMPR2, and found that this antibody is also capable of neutralizing the ligand binding function of BMPR2-ECD. We extended the results by examining the ability of 3F6 to block endogenous BMPR2 function in the BMP-responsive HEK293T (human kidney embryonic 293 translation) cell line. Consistent with the results of our cell-free system, pre-treatment of HEK293T cells with 3F6 leads to reduced sensitivity to in response to BMP pathway activation by BMP2. These results provide proof-of-concept data for future studies evaluating inhibition of BMPR2 function in vivo as a means to reduce age-related bone loss.



Conclusions:

Our results provide proof-of-concept data that BMPR2 function can be blocked using a neutralizing antibody approach. Specifically, the mouse monoclonal antibodies 3F6 and 1F12 block BMPR2-ECD ligand-binding in a cell-free immunoprecipitation assay. 3F6 was additionally shown to block BMP2-induced pathway activation in HEK293T cells. An incidental finding from our study is that BMPR2 is the major receptor for BMP2 in HEK293T cells.

Current and Future Directions:

- 1) Examination of 3F6 and 1F12 neutralizing activity of BMPR2 function in skeletal cells in vitro.
- 2) Examination of systemic delivery of 3F6 and/or 1F12 in vivo as a means of regulating postnatal bone mass.

Acknowledgements:

This work was performed in collaboration with:
Laura Schoerning
Aaron Hudnall

Jordan Newby
Julia Hum, PhD

Warren Lawless

Supported by a MU-COM Faculty Research Development award issued to JWL.

Follow this link for a video presentation of this poster and to leave feedback:



<http://www.jonlowery.com/presentations>

Follow this link to the Lowery Lab Website:



<http://www.jonlowery.com/research>